

Product Documentation

Akoline PG7™

8691

Version

Date 2024 02 06

To whom it may concern

Dear valued customer:

The purpose of this document is to provide you with the information required to evaluate the safety of this product to fulfil the legal requirements. The second purpose of the document is to provide you with all information required during the coding process. AAK has gathered the questions received throughout the years and collected the answers within this document. The document is strictly addressing the cosmetic and personal care applications, thus having no intention to cover, pharmaceutical, food or other applications. As the regulatory requirements increases on the answers given as well as the number of questionnaires increases, AAK has chosen to focus on quality and to give you an answer within a reasonable time. This document represents the answer to your questionnaire. AAK has tried to be as complete and accurate as possible in providing the information and feels comfortable it covers the needs for you. In the case AAK does not possess data or information for a particular subject it is stated in the document.

A handwritten signature in blue ink, appearing to read 'Staffan Norberg', followed by a large, stylized blue checkmark or 'L' shape.

Head of Development AAK-PC

Staffan Norberg

Contents

1.	IDENTIFICATION	8
1.1	Identification	
2.	CHEMICAL AND PHYSICAL DATA	9
2.1	Specifications	
2.2	Typical values	
2.3	Certificate of Analysis	
2.4	Auxiliary chemical and physical data	
3.	RAW MATERIAL	10
3.1	Biological data	
3.2	Composition breakdown	
4.	PRODUCTION	11
4.1	Production data	
5.	BY-PRODUCTS AND OTHER IMPURITIES	12
5.1	AAK Contaminant standard	
5.2	Other impurities specific substances	
5.3	Impurities general statements	
5.3.1	Allergenes	
5.3.2	Proteins	
5.3.3	VOC Volatile Organic Compounds	
5.3.4	Sulphonates	
5.3.5	Parabens	
5.3.6	Phthalates	
5.3.7	Silicones	
6.	STABILITY DATA	13
6.1	Stability Data	
7.	HUMAN HEALTH HAZARD ASSESSMENT	14

7.1	General read-across consideration and justification
7.2	Acute toxicity
7.2.1	Acute oral toxicity
7.2.2	Acute inhalation toxicity
7.2.3	Acute dermal toxicity
7.2.4	Acute toxicity by other exposure routes
7.2.5	Summary and discussion of acute toxicity
7.3	Irritation & corrosivity
7.3.1	Skin irritation and corrosivity
7.3.2	Eye & mucous membrane irritation and corrosivity
7.3.3	Summary and discussion on irritation and corrosivity
7.4	Skin sensitization
7.4.1	Summary and discussion of sensitisation
7.5	Repeated dose, sub-chronic and chronic toxicity
7.5.1	Oral administration
7.5.2	Inhalation studies
7.5.3	Dermal administration
7.5.4	Other routes of administration
7.5.5	Human information
7.6	Reproduction toxicity
7.6.1	Non-human information
7.6.2.	Human information
7.6.3	Developmental toxicity/teratogenicity
7.6.4	Summary and discussion of reproductive toxicity
7.7	Mutagenicity/genotoxicity
7.7.1	In vitro data
7.7.2	In vivo data
7.7.3	Human information

7.7.4	Summary and discussion of mutagenicity	
7.8	Carcinogenicity	
7.8.1	Non-human information	
7.8.2	Human information	
7.8.3	Summary and discussion of carcinogenicity	
7.9	Toxicokinetics (absorption, metabolism, distribution and elimination (ADME))	
7.9.1	Oral administration	
7.9.2	Dermal administration	
7.9.3	Inhalation route	
7.10	Photoinduced toxicity	
7.10.1	Phototoxicity: photoirritation / photosensitisation	
7.10.2	Phototoxicity: photomutagenicity / photoclastogenicity	
7.10.3	Other relevant human studies (clinical)	
7.11	Special investigations	
7.12	Summary and NOAEL statement	
8.	ECOLOGICAL DATA	35
8.1	Degradability	
8.2	Accumulation	
8.3	Aquatic toxicity	
9.	REGULATORY	40
9.1	EU	
9.1.1	EU Cosmetic Regulation EC 1223/2009	
9.1.2	EU Cosmetic Regulation EC 1223/2009, Annex II and III	
9.1.3	EU REACH 1907/2006	
9.1.4	EU SVHC (Substance of Very High Concern)	
9.1.5	Other	
9.2	USA	
9.2.1	US (California) Proposition 65	

9.2.2	China – NMPA	
9.2.3	UK REACH	
9.2.4	Turkey – KKDIK	
9.2.5	Australia – TGA	
9.2.6	Other	
9.3	Other non-Country specific regulatory issues	
9.3.1	Animal testing	
9.3.2	Nano particlesTurkey	
9.3.3	Nagoya Protocol / Biodiversity and Access Benefit Sharing regulation	
9.3.4	CITES	
9.3.5	CMR	
9.3.6	Other	
9.4	Inventory lists	
10	General statements and standards	42
10.1	Official standards	
10.2	Private standards	
10.2.1	Ecocert, Cosmos or Natrue	
10.2.2	Vegan and Vegetarian claim	
10.2.3	Other	
10.3	Other Statements	
10.3.1	BSE/TSE statements	
10.3.2	GMO statement	
10.3.3	Other	
11.	CERTIFICATES	43
11.1	HALAL	
11.2	Kosher	

11.3	ISO 9001	
11.4	EFfCI GMP	
11.5	Food Safety/ FSSC 22000	
11.6	Other	
12.	PATENTS	44
12.1	Patents	
13.	TRANSPORTS AND HANDLING	45
13.1	Transports	
13.2	storage unopen package	
13.3	Handling of product for use	
13.3.1	Use of full package	
13.3.2	Use of full package for partly use	
14.	REFERENCES	46
14.1	References	
15.	DISCLAIMER	47
15.1	Disclaimer	
16.	APPENDIX	
C0043	Certificate of Analasys	48
T0005	Process flowchart	49

1.1 Identification

Producer: AAK Sweden AB, Västra kajen SE-374 82 Karlshamn, Sweden
Tradename: Akoline PG7™
Art. No: 8691
Country of Origin EU

This product is used globally. As the product may fit in the definition of several CAS numbers, AAK give examples of alternative CAS number to be used for instance in inventory lists search.

	INCI	CAS Number	EC number
EU /AAK first choice	Polyglyceryl-3-Stearate	27321-72-8	248-403-2
US	Polyglyceryl-3-Stearate	27321-72-8	248-403-2
China*	聚甘油-3 硬脂酸酯	27321-72-8	248-403-2
Alternative INCI	Polyglyceryl-3-Stearate	26855-43-6	248-403-2

*) For NMPA information see section 9.2.2 China – NMPA



Margrét Viborg
Global Regulatory Affairs Manager

2.1 Specifications

For specification see Product Data Sheet (PDS)

Download latest version at www.aakpersonalcare.com/

2.2 Typical values

For typical values see Product Data Sheet (PDS)

Download latest version at www.aakpersonalcare.com/

2.3 Certificate of Analysis

For example of COA, see Appendix.

2.4 Auxiliary chemical and physical data

Molecular weight ~880 g/mol

3.1 Biological data

Botanical origin

INCI	Botanical origin	^{*)} Geographical origin	Part used	Content %	Wild grown or cultivated
Polyglyceryl-3-Stearate	Elaeis Guineensis	Malaysia or Indonesia	Fruit flesh	100	Cultivated

^{*)}Geographical origin may change

3.2 Composition breakdown

INCI name (EU)	CAS	EINECS	Average Content %	Function
Polyglyceryl-3-Stearate	27321-72-8	248-403-2	100	Emulsifier

Palm content:

☒ Containing palm

☐ RSPO SG:

☒ RSPO MB: CU-RSPO SCC-817671

☐ Do not contain Palm



Margrét Viborg
Global Regulatory Affairs Manager
Personal Care, AAK Sweden AB

4.1 Production data

The following operations are used in the processing of this ingredient

Process		Comment
Mechanical extraction		
Solvent extraction		
Refining		
Deodorising		
Hydrogenation		
Interesterification		
Esterification	X	
Winterisation		
Solvent Fractionation		
Dry Fractionation		
Ethoxylation		
Molecular distillation		
Other processing	X	See attached flowchart

5. BY-PRODUCTS AND OTHER IMPURITIES

5.1 AAK Contaminant standard

Not available, for more information please contact AAK.

5.2 Other Impurities specific substances

Download latest version of “AAK personal Care position on impurities” at aakpersonalcare.com

5.3 Impurities AAK Cosmetic Products

5.3.1 Allergens

Download “General statements AAK Cosmetic Ingredients” at aakpersonalcare.com

5.3.2 Proteins

Download “General statements AAK Cosmetic Ingredients” at aakpersonalcare.com

5.3.3 VOC – Volatile Organic Compounds

Download “General statements AAK Cosmetic Ingredients” at aakpersonalcare.com

5.3.4 Sulphonates

Download “General statements AAK Cosmetic Ingredients” at aakpersonalcare.com

5.3.5 Parabens

Download “General statements AAK Cosmetic Ingredients” at aakpersonalcare.com

5.3.6 Phthalates

Download “General statements AAK Cosmetic Ingredients” at aakpersonalcare.com

5.3.7 Silicones

Download “General statements AAK Cosmetic Ingredients” at aakpersonalcare.com



6.1 Stability data

OSI Value @ 120C No data

Storage @ 20C

Peroxide value 12 month: No data 24 month: No data

Storage @ 40C

Peroxide value 12 month: No data 24 month: No data

7 Human Health and Environmental Hazard Assessment

Akoline PG7

7.01 General read-across consideration and justification

Test name:

CIR Safety assessment

Method and laboratory:

Safety assessment and review of various polyglycerol ester based cosmetic ingredients

Test material:

274 polyglycerol esters with 2 to 20 glycerol units and a variety of fatty acids

Results:

The CIR panel concludes that the polyglycerol esters are safe in cosmetics in the current practices and when formulated to be non-irritating.

Read across

Statement

Reference ID:

S-345 Final Report: Safety assessment of polyglyceryl fatty acid esters as used in cosmetics. Cosmetic Ingredient Review, November 14, 2016. Available at <https://www.cir-safety.org/>

Test name:

EFSA scientific opinion and safety review

Method and laboratory:

EFSA scientific opinion and safety review on polyglycerol polyricinoleate (E 476)

Test material:

Polyglycerol polyricinoleate

Results:

EFSA concludes that polyglycerol polyricinoleate is not of concern for acute or long-term toxicity, including reproductive toxicity, genotoxicity and carcinogenicity. The oral/systemic ADI (acceptable daily intake) is set to 25 mg/kg bw.

Read across

Statement

Reference ID:

S-346 Re-evaluation of polyglycerol polyricinoleate (E 476) as a food additive. EFSA Journal 15(3): 4743, 54pp, (2017)

Test name:

EFSA scientific opinion and safety review

Method and laboratory:

EFSA scientific opinion and safety review on polyglycerol esters of fatty acids (E 475)

Test material:

Polyglycerol esters of fatty acids

Results:

EFSA concludes that polyglycerol esters of fatty acids are not of concern for acute or long-term toxicity, including reproductive toxicity, genotoxicity and carcinogenicity. There is no need to set an ADI (acceptable daily intake) for food applications.

Read across

Statement

Reference ID:

S-347 Re-evaluation of polyglycerol esters of fatty acids (E 475) as a food additive. EFSA Journal 15(12): 5089, 32pp, (2017)

Test name:

Summary and review of polyglyceryl polyricinoleate

Method and laboratory:

Review and summary of studies on polyglycerol polyricinoleate

Test material:

Polyglycerol polyricinoleate

Results:

Review of preparation, use and biological studies on polyglycerol polyricinoleate

Comments:

General background information

Read across

Statement

Reference ID:

S-348 Overview of the preparation, use and biological studies on polyglycerol polyricinoleate (PGPR). Wilson, R., van Schie, B.J. & Howes, D., Food Chemical Toxicology, 36, 711-718, (1998)

Test name:

REACH registration dossier

Method and laboratory:

REACH registration dossier with toxicological and environmental data

Test material:

Polyglycerol-3

Read across

Statement

Reference ID:

S-352 ECHA, REACH registration 'Reaction mass of 3-[3-(2,3-dihydroxypropoxy)-2-hydroxypropoxylpropane-1,2-diol...',
<https://echa.europa.eu/registration-dossier/-/registered-dossier/13319>
Accessed 2020-08-18

Test name:

REACH registration dossier

Method and laboratory:

REACH registration dossier with toxicological and environmental data

Test material:

Polyglycerol-3 caprate/laurate

Read across

Statement

Reference ID:

S-353 ECHA, REACH registration 'Reaction products of decanoic acid and lauric acid with glycerol and polyglycerol',

<https://echa.europa.eu/registration-dossier/-/registered-dossier/25834>

Accessed 2020-08-18

Test name:

REACH registration dossier

Method and laboratory:

REACH registration dossier with toxicological and environmental data

Test material:

Polyglycerol-3 caprate

Read across

Statement

Reference ID:

S-354 ECHA, REACH registration '1,2,3-propantriol, homopolymer, decanoate', CAS 74504-65-7

<https://echa.europa.eu/registration-dossier/-/registered-dossier/23621>

Accessed 2020-08-18

Test name:

REACH registration dossier

Method and laboratory:

REACH registration dossier with toxicological and environmental data

Test material:

Polyglycerol-3

Read across

Statement

Reference ID:

S-355 ECHA, REACH registration '1,2,3-propantriol, homopolymer', CAS 25618-55-7

<https://echa.europa.eu/registration-dossier/-/registered-dossier/14285>

Accessed 2020-08-18

Polyglycerol esters are a large group of emulsifiers and viscosity regulating substances that comprise a polar group made from polyglycerol and a non-polar group consisting of fatty acids. The majority of the polyglycerol esters are based on polyglycerol-3, which are mixtures of di-, tri-, and tetraglycerol with varying amounts of glycerol and higher oligomers. A typical composition is quoted in the EFSA review (S-347) as 1.3% glycol, 26.2% glycerol, 4.9% cyclic diglycerol, 22.7% diglycerol, 15.6% triglycerol, 9.1% tetraglycerol, 6.0% pentaglycerol, 3.8% hexaglycerol and 3.6% higher polyglycerols. Another group of common polyglyceryl esters are based on polyglycerol-10, with a distribution of oligomers centering on 10 glycerol units.

The polyglycerol may be esterified to one or more fatty acids or mixtures of fatty acids. The saturated fatty acids range from caprylic (C8) to behenic (C22), and unsaturated esters are normally based on oleic and/or linoleic acids (C18:1, C18:2). Iso-esters with branched fatty acids are also available (mainly isostearamides). Hydroxy acids such as ricinoleic acid and dihydroxystearate are also used, the ricinoleic acid normally as a polymer of ricinoleic acid ("polyricinoleates"). Many commercial ingredients are named after the starting raw material, for example "coconut oil polyglyceryl-6 esters". Such ingredients are characterized by the fatty acids found in the starting materials so for the coconut based material mainly lauric/myristic/palmitic/stearic acids.

The polyglycerols are esterified with one or more fatty acids. For example, polyglyceryl-3 stearate has on average one fatty acid per polyglycerol but the actual composition will be a mixture of mono-esters, di-esters, tri-esters etc and unreacted polyglycerol.

The toxicology and environmental properties of the polyglyceryl esters depend on the properties of the substances themselves but also on the metabolites (polyglycerols and fatty acids). For this reason, read-across can be made over a wide range of different polyglyceryl esters. In this review both saturated, unsaturated, branched and hydroxy-acid based materials are considered equal from a toxicology and environmental point of view and extensive read-across is applied. The main point for caution is the skin and eye irritation potential of the shorter chain (C8-C12) esters, due to their high surface activity, which will differ from the longer chain alternatives.



7.02 Acute toxicity

7.02.1 Acute oral toxicity

Test name:

Summary of acute toxicity of polyglycerol esters

Method and laboratory:

Review and summary of several studies on the acute toxicity of polyglyceryl esters

Test material:

Polyglycerol esters of various saturated, unsaturated and branched fatty acids

Results:

LD50(oral): >5g/kg bw (rat)

LD50(dermal): >5g/kg bw (rat)

Comments:

(Table 12 in report).

Read across

Read across Similar chemical structures

Reference ID:

S-345 Final Report: Safety assessment of polyglyceryl fatty acid esters as used in cosmetics. Cosmetic Ingredient Review, November 14, 2016. Available at <https://www.cir-safety.org/>

Test name:

Acute oral toxicity

Method and laboratory:

Summary of acute oral toxicity data in EFSA review on polyglycerol polyricinoleate (PGPR).

Test material:

Polyglycerol polyricinoleate

Ricinoleic acid

Castor oil

Polyglycerols

Results:

LD50(rat, PGPR): >20000 mg/kg bw

LD50(mouse, PGPR): >100000 mg/kg bw

LD50(rat, polyglycerols): >2000 mg/kg bw

Comments:

Castor oil and ricinoleic acid cause diarrhoea and potential disturbance of the intestinal mucosa.

Read across

Read across Similar chemical structures

Reference ID:

S-346 Re-evaluation of polyglycerol polyricinoleate (E 476) as a food additive. EFSA Journal 15(3): 4743, 54pp, (2017)

7.02.2 Acute inhalation toxicity

No actual tests have been carried out and literature data has not been found for this chapter.

7.02.3 Acute dermal toxicity

Test name:

Summary of acute toxicity of polyglycerol esters

Method and laboratory:

Review and summary of several studies on the acute toxicity of polyglyceryl esters

Test material:

Polyglycerol esters of various saturated, unsaturated and branched fatty acids

Results:

LD50(oral): >5g/kg bw (rat)

LD50(dermal): >5g/kg bw (rat)

Comments:

(Table 12 in report).

Read across

Read across Similar chemical structures

Reference ID:

S-345 Final Report: Safety assessment of polyglyceryl fatty acid esters as used in cosmetics. Cosmetic Ingredient Review, November 14, 2016. Available at <https://www.cir-safety.org/>

7.02.4 Acute toxicity by other exposure routes

No actual tests have been carried out and literature data has not been found for this chapter.

7.02.5 Summary and discussion of acute toxicity

Polyglycerol esters of fatty acids, including 'polyglycerol-3 stearate' and 'polyglycerol-3 polyricinoleate', have a very long history of safe use in a wide range of nutritional (food and feed), cosmetic and industrial applications. The acute oral, inhalation or dermal toxicity is low and therefore not considered to pose an issue for human health under normal and foreseeable handling and use conditions. There is no Acceptable Daily Intake (ADI) assigned to polyglycerol esters of saturated fatty acids. The ADI for polyglycerol polyricinoleate is set to 25 mg/kg bw/day.

7.03 Irritation & corrosivity

7.03.1 Skin irritation and corrosivity

Test name:

Skin irritation and sensitization

Method and laboratory:

Review and summary of skin irritation and sensitization for polyglycerol esters (Table 14 in CIR report). 7 alternative studies, 36 animal studies and 24 human studies are reported.

Test material:

Saturated, unsaturated and branched polyglycerol esters

Results:

Polyglycerol esters are not irritating to the skin
Polyglycerol esters are not sensitizing

Comments:

A small number of reports indicate slight to moderate irritation (5-6 out of 67 reports)

Read across

Read across Similar chemical structures

Reference ID:

S-345 Final Report: Safety assessment of polyglyceryl fatty acid esters as used in cosmetics. Cosmetic Ingredient Review, November 14, 2016. Available at <https://www.cir-safety.org/>

7.03.2 Eye & mucous membrane irritation and corrosivity

Test name:

Eye irritation

Method and laboratory:

Review and summary of eye irritation for polyglycerol esters (Table 15 in CIR report). 23 alternative studies and animal studies and 1 human study are reported.

Test material:

Saturated, unsaturated and branched polyglycerol esters

Results:

Polyglycerol esters are not irritating to minimally irritating to the eye.

Read across

Read across Similar chemical structures

Reference ID:

S-345 Final Report: Safety assessment of polyglyceryl fatty acid esters as used in cosmetics. Cosmetic Ingredient Review, November 14, 2016. Available at <https://www.cir-safety.org/>

7.03.3 Summary and discussion on irritation and corrosivity

Based on the safety assessments and reviews it is concluded that polyglycerol esters of saturated, unsaturated, branched and hydroxy fatty acids pose a minimal risk to cause skin or eye irritation in normal use concentrations. It is also concluded that the polyglycerol esters of fatty acids are not corrosive to skin, eyes or mucous membranes.

7.04 Skin sensitization

Test name:

Skin irritation and sensitization

Method and laboratory:

Review and summary of skin irritation and sensitization for polyglycerol esters (Table 14 in CIR report). 7 alternative studies, 36 animal studies and 24 human studies are reported.

Test material:

Saturated, unsaturated and branched polyglycerol esters

Results:

Polyglycerol esters are not irritating to the skin

Polyglycerol esters are not sensitizing

Comments:

A small number of reports indicate slight to moderate irritation (5-6 out of 67 reports)

Read across

Read across Similar chemical structures

Reference ID:

S-345 Final Report: Safety assessment of polyglyceryl fatty acid esters as used in cosmetics. Cosmetic Ingredient Review, November 14, 2016. Available at <https://www.cir-safety.org/>

7.04.1 Summary and discussion of sensitization

According to read-across studies, presented in the CIR safety report and in REACH registration dossiers, polyglycerol esters such as 'polyglycerol-3 stearate' and 'polyglycerol-3 polyricinoleate' are not sensitizing.

7.05 Repeated dose, sub-chronic and chronic toxicity
7.05.1 Oral administration

Test name:

Summary of sub-chronic toxicity of polyglycerol esters

Method and laboratory:

Review and summary of the sub-chronic toxicity of polyglyceryl esters in rat (10M/10F Sprague-Dawley rats, duration 90 days, dosage 2.5, 5.0 and 10.0% in diet)

Test material:

Polyglycerol-10 decaoleate

Results:

No treatment related adverse effects were detected.

Comments:

Also cited in EFSA review (S-347)

Read across

Read across Similar chemical structures

Reference ID:

S-345 Final Report: Safety assessment of polyglyceryl fatty acid esters as used in cosmetics. Cosmetic Ingredient Review, November 14, 2016. Available at <https://www.cir-safety.org/>

Test name:

Summary of chronic toxicity of polyglycerol esters

Method and laboratory:

Review and summary of the chronic toxicity of polyglyceryl esters in mice (25M/25F mice, duration 80 weeks, dosage 5% in diet) and rats (28M/28F rats, duration 2 years, dosage 5% in diet).

Test material:

Polyglycerol esters

Results:

No treatment related adverse effects were detected. Liver and kidney weights in female mice were higher than in control group (5% groundnut oil).

Comments:

Also cited in EFSA review (S-347)

Read across

Read across Similar chemical structures

Reference ID:

S-345 Final Report: Safety assessment of polyglyceryl fatty acid esters as used in cosmetics. Cosmetic Ingredient Review, November 14, 2016. Available at <https://www.cir-safety.org/>

Test name:

Sub-chronic and short-term oral toxicity

Method and laboratory:

Summary of short-term oral toxicity data in EFSA review on polyglycerol polyricinoleate (PGPR).

- 1) Mouse 100M/100F, duration 14 days, dosage 0.5-8% PGPR in diet (450-7200 mg/kg bw/day).
- 2) Mouse 140M/140F, duration 14 days, dosage 0, 5, 10 & 15% in diet (0, 4500, 9000 and 13500 mg/kg bw/day).
- 3) Rat 6M/6F, duration 5 days, dosage 10 ml/kg bw/day by gavage (10000 mg/kg bw/day).
- 4) Rat 24M/24F, duration 14 days, dosage 18% in diet (16200 mg/kg bw/day).

Test material:

Polyglycerol polyricinoleate

Results:

- 1) No adverse effects observed up to 1800 mg/kg bw/day. Increased liver weights observed at higher dosages.
- 2) Comparable growth rates between treatment and control groups. Increased liver weights in PGPR groups.
- 3) No adverse effects observed.
- 4) Increased liver weights seen in PGPR groups

The EFSA panel concludes that the increased liver weight is reversible and due to high levels of the test substance.

Comments:

2 more studies with similar conditions and outcomes reported in the EFSA review (S-346).

Read across

Read across Similar chemical structures

Reference ID:

S-346 Re-evaluation of polyglycerol polyricinoleate (E 476) as a food additive. EFSA Journal 15(3): 4743, 54pp, (2017)

Test name:

Chronic and carcinogenicity toxicity

Method and laboratory:

Summary of long-term and carcinogenicity toxicity data in EFSA review on polyglycerol polyricinoleate (PGPR).

1) Mice 100M/100F in four treatment groups, duration 80 weeks, dosage 5% in diet (7500 mg/kg bw/day).

2) Rats 48M/48F, duration 30 and 45 weeks (2 studies), dosage 9% in diet (4500 mg/kg bw/day).

3) Rats 30M/30F, duration 2 years, dosage 5% (2500 mg/kg bw/day).

Test material:

Polyglycerol polyricinoleate

Results:

1) No treatment related adverse effects observed. No increased incidence of tumors observed. NOAEL (mice) = 7500 mg/kg bw/day.

2) No treatment related adverse effects were observed. NOAEL (rat) = 4500 mg/kg bw/day.

3) No treatment related adverse effects observed. No increased incidence of tumors observed. NOAEL (rat) = 2500 mg/kg bw/day.

Read across

Read across Similar chemical structures

Reference ID:

S-346 Re-evaluation of polyglycerol polyricinoleate (E 476) as a food additive. EFSA Journal 15(3): 4743, 54pp, (2017)

Test name:

Chronic toxicity and carcinogenicity

Method and laboratory:

Summary of chronic toxicity and carcinogenicity studies in EFSA review on polyglycerol esters

1) mice, 25M/25F, duration 80 weeks, dosage 5% in diet (7500 mg/kg bw/day).

2) rats, 28M/28F per dosage group, duration 2 years, dosage 5% in diet (2500 mg/kg bw/day)

Test material:

Polyglycerol esters of fatty acids

Results:

1) No treatment related adverse effects were observed. No increased incidence of tumors. NOAEL (mice) = 7500 mg/kg bw/day.

2. No treatment related adverse effects were observed. No increased incidence of tumors. NOAEL (rat) = 2500 mg/kg bw/day.

Read across

Read across Similar chemical structures

Reference ID:

S-347 Re-evaluation of polyglycerol esters of fatty acids (E 475) as a food additive. EFSA Journal 15(12): 5089, 32pp, (2017)

7.05.2 Inhalation studies

No actual tests have been carried out and literature data has not been found for this chapter.

7.05.3 Dermal administration

No actual tests have been carried out and literature data has not been found for this chapter.

7.05.4 Other routes of administration

No actual tests have been carried out and literature data has not been found for this chapter.

7.05.5 Human studies

7.05.6 Summary and discussion

The highest oral NOAEL for polyglycerol esters of fatty acids based on the studies reported above, could be considered to be 5% in feed, equivalent to an estimated 2500 mg/kg bw/day.

Polyglycerol-3 stearate, polyglycerol-3 ricinoleate and other substances from the same read-across category present low systemic toxicity upon repeated dose oral exposure, and the repeated dose dermal toxicity is also expected to be minimal. The substances are metabolized into glycerol, polyglycerols and fatty acids. The glycerol and the fatty acids are metabolized according to well established pathways. Polyglycerol is excreted without metabolism in the urine or faeces. Furthermore, being non-volatile, repeated inhalation exposure is not considered to pose an issue for human health under normal and foreseeable handling and use conditions.

Based on the above information, the substance does not qualify for repeated dose toxicity classification according to Directive 67/548/EC or Regulation 1272/2008/EC.

7.06 Reproduction toxicity

7.06.1 Non-human studies

Test name:

Reproduction study PGPR

Method and laboratory:

Three-generation reproduction study in rats.

Test material:

Polyglycerol polyricinoleate (ADMUL WOL), 1.5% in diet

Results:

No adverse, treatment related, effects on the reproductive function were observed in this study.

Comments:

Also cited in EFSA review (S-346). Study of limited value due to methodological problems.

Read across

Read across Similar chemical structures

Reference ID:

S-350 A Three-generation reproduction study on polyglycerol polyricinoleate (PGPR) in Wistar rats. Wilson, R. & Smith, M., Food Chemical Toxicology, 36, 739-741, (1998)

Test name:

Reproduction and development toxicity

Method and laboratory:

Review and summary of reproduction and development toxicity of polyglyceryl esters in rats (OECD Guideline 422, dosage 0, 100, 300 and 1000 mg/kg bw/day).

Test material:

Polyglycerol-3 diisostearate

Results:

NOAEL for systemic effects was determined to be >1000mg/kg bw/day. No treatment related adverse effects were detected.

Comments:

Also cited in EFSA review (S-347)

Read across

Read across Similar chemical structures

Reference ID:

S-345 Final Report: Safety assessment of polyglyceryl fatty acid esters as used in cosmetics. Cosmetic Ingredient Review, November 14, 2016. Available at <https://www.cir-safety.org/>

7.06.2 Human studies

No actual tests have been carried out and literature data has not been found for this chapter.

7.06.3 Developmental toxicity/teratogenicity

No actual tests have been carried out and literature data has not been found for this chapter.

7.06.3.1 Non-human studies

No actual tests have been carried out and literature data has not been found for this chapter.

7.06.3.2 Human studies

No actual tests have been carried out and literature data has not been found for this chapter.

7.06.4 Summary and discussion of reproductive toxicity

No reliable published studies on reproductive and developmental toxicity for polyglycerol esters of fatty acids or other substances in the same read-across category have been found. One study on polyglycerol polyricinoleate is available but the study is not well designed and lack important details in the documentation. A study on polyglycerol-3 diisostearate is reported in the CIR review (S-345) showing no effects on reproduction up to the highest tested concentration (1000 mg/kg bw/day). Several problems relating to the execution of the study are reported and the validity of the study is questionable.

7.07 Mutagenicity/genotoxicity

7.07.1 In vitro data

Test name:

Mutagenicity and genotoxicity

Method and laboratory:

Review and summary of mutagenicity and genotoxicity polyglycerol esters (Table 13 in CIR report). 27 in vitro tests are reported (Ames test, mammalian cell mutation assays, chromosomal aberration assays, with and without metabolic activation)

Test material:

Saturated, unsaturated and branched polyglycerol esters

Results:

The tested polyglycerol esters are not mutagenic, genotoxic or clastogenic.

Read across

Read across Similar chemical structures

Reference ID:

S-345 Final Report: Safety assessment of polyglyceryl fatty acid esters as used in cosmetics. Cosmetic Ingredient Review, November 14, 2016. Available at <https://www.cir-safety.org/>

7.07.2 In vivo data

No actual tests have been carried out and literature data has not been found for this chapter.

7.07.3 Human studies

No actual tests have been carried out and literature data has not been found for this chapter.

7.07.4 Summary and discussion of mutagenicity

Based on in vitro studies on gene mutation tests in bacteria and mammalian cells as well as mammalian chromosome aberration tests on substances that are structurally relevant, it is concluded that polyglycerol esters of fatty acids do not present any concern for mutagenicity and genotoxicity.

7.08 Carcinogenicity

7.08.1 Non-human studies

Test name:

Carcinogenic properties

Method and laboratory:

Assessment of carcinogenic potential for PGPR in rat & mouse.

Rat: 60 M/ 60 F, duration 2 years, dosage 5% in diet

Mice: 25 M/ 25F, duration 80 weeks, dosage 5% in diet

Test material:

Polyglycerol polyricinoleate
(ADMUL WOL)

Results:

No carcinogenic effects were detected. No adverse effects on growth, food consumption, longevity or haematology. No treatment related histological changes. Increase in liver and kidney weight was observed in both mice and rats.

Read across

Read across Similar chemical structures

Reference ID:

S-349 Assessment of the carcinogenic potential of polyglycerol polyricinoleate (PGPR) in rats and mice, Smith, M.R., Wilson, R. & Hepburn, P.A., Food Chemical Toxicology, 36, 747-754, (1998)

Test name:

Chronic and carcinogenicity toxicity

Method and laboratory:

Summary of long-term and carcinogenicity toxicity data in EFSA review on polyglycerol polyricinoleate (PGPR).

1) Mice 100M/100F in four treatment groups, duration 80 weeks, dosage 5% in diet (7500 mg/kg bw/day).

2) Rats 48M/48F, duration 30 and 45 weeks (2 studies), dosage 9% in diet (4500 mg/kg bw/day).

3) Rats 30M/30F, duration 2 years, dosage 5% (2500 mg/kg bw/day).

Test material:

Polyglycerol polyricinoleate

Results:

1) No treatment related adverse effects observed. No increased incidence of tumors observed. NOAEL (mice) = 7500 mg/kg bw/day.

2) No treatment related adverse effects were observed. NOAEL (rat) = 4500 mg/kg bw/day.

3) No treatment related adverse effects observed. No increased incidence of tumors observed. NOAEL (rat) = 2500 mg/kg bw/day.

Read across

Read across Similar chemical structures

Reference ID:

S-346 Re-evaluation of polyglycerol polyricinoleate (E 476) as a food additive. EFSA Journal 15(3): 4743, 54pp, (2017)

7.08.2 Human studies

No actual tests have been carried out and literature data has not been found for this chapter.

7.08.3 Summary and discussion of carcinogenicity

Several chronic toxicity and carcinogenicity studies are summarized in the EFSA reviews on polyglyceryl esters and polyglyceryl polyricinoleate (S-346 & S-347). The studies show that no increases in tumor incidence can be seen in mice fed 7500 mg/kg bw/day or rats fed 2500 mg/kg bw/day. A NOAEL of 2500 mg/kg bw/day is derived from these studies.

It is concluded that polyglycerol esters are unlikely to be a health hazard from a carcinogenicity point of view.

7.09 Toxicokinetics: absorption, metabolism, distribution and elimination (ADME)

7.09.1 Oral administration

Test name:

Toxicokinetics (ADME)

Method and laboratory:

The absorption, tissue distribution, metabolism and excretion of radiolabelled polyglycerol polyricinoleate was studied in rats in several ways. Both radiolabelled polyglycerol and radiolabelled fatty acids were used. The digestion of polyglycerol polyricinoleate by pancreatic lipase was also studied in vitro.

Test material:

Polyglycerol polyricinoleate

Results:

Polyglycerol esters are completely hydrolysed with the aid of lipases to the corresponding polyglycerol and fatty acids. Lower polyglycerols are excreted in the urine while higher polyglycerols are bound in the faeces. Glycerol and fatty acids are metabolized in the normal manner.

Read across

Read across Similar chemical structures

Reference ID:

S-351 The fate of ingested glyceran esters of condensed castor oil fatty acids [Polyglycerol polyricinoleate (PGPR)] in the rat. Howes, D., Wilson, R. & James, C.T., Food Chemical Toxicology, 36, 719-738, (1998)

Test name:

Summary of adsorption, distribution, metabolism and excretion of polyglycerol esters

Method and laboratory:

Review and summary of several studies on the ADME of polyglyceryl esters

Test material:

Polyglycerol esters of stearic, oleic and linoleic acids

Results:

Polyglycerol esters are hydrolyzed by lipases into polyglycerol(s) and fatty acids. The fatty acids are metabolized in the normal manner while the polyglycerols are excreted.

Read across

Read across Similar chemical structures

Reference ID:

S-345 Final Report: Safety assessment of polyglyceryl fatty acid esters as used in cosmetics. Cosmetic Ingredient Review, November 14, 2016. Available at <https://www.cir-safety.org/>

7.09.2 Dermal administration

No actual tests have been carried out and literature data has not been found for this chapter.

7.09.3 Inhalation route

No actual tests have been carried out and literature data has not been found for this chapter.

7.10 Photoinduced toxicity

7.10.1 Phototoxicity: photoirritation / photosensitisation

Test name:

Phototoxicity and photosensitization

Method and laboratory:

Review and summary of phototoxicity (guinea pigs, 10F, 0.5 ml polyglycerol-10 nonaisostearate, 150 mJ/cm² UVB, 7000 mJ/cm² UVA) and photosensitization (guinea pig, n=11, 7000 mJ/cm² UVA, induction with 2 intradermal injections, challenge after 16 days, application of test substance + irradiation with 7 mJ/cm² UVA)

Test material:

Polyglycerol-10 nonaisostearate

Results:

Polyglycerol-10 nonaisostearate is not phototoxic or a photosensitizer.

Read across

Read across Similar chemical structures

Reference ID:

S-345 Final Report: Safety assessment of polyglyceryl fatty acid esters as used in cosmetics. Cosmetic Ingredient Review, November 14, 2016. Available at <https://www.cir-safety.org/>

7.10.2 Phototoxicity: photomutagenicity / photoclastogenicity

No actual tests have been carried out and literature data has not been found for this chapter.

7.10.3 Other relevant human studies (clinical)

No actual tests have been carried out and literature data has not been found for this chapter.

7.11 Special investigations

No actual tests have been carried out and literature data has not been found for this chapter.

7.12 Summary and NOAEL statement

Based on the data presented in chapter 7.1-7.11 a NOAEL value of 2500 mg/kg bw/day for systemic exposure is given for polyglycerol esters of fatty acids such as 'polyglycerol-3 stearate' and 'polyglycerol-3 ricinoleate'.

8 Ecological data

8.01 Degradability

Test name:

Biodegradation

Method and laboratory:

Biodegradation was followed by measuring BOD/ThOD (biological oxygen demand/theoretical oxygen demand) over 28 days in closed bottles using inoculum from a municipal sewage plant.

Test material:

Diglycerol Triglycerol

Tetraglycerol

Diglyceryl esters from caprylate to laurate

Triglyceryl esters from caprylate to laurate

Results:

All tested polyglycerols and polyglyceryl esters were readily biodegradable.

Read across

Read across Similar chemical structures

Reference ID:

S-324 Enzymatic synthesis, surface activity, antimicrobial properties and biodegradability of di- and triglycerol fatty acid esters, J Jpn Oil Soc, 48(7), 681-725, (1999)

Test name:

Biodegradability OECD 301F

Method and laboratory:

OECD 301F Manometric Respirometry Test 1992

Aerobic biodegradability of organic compounds. 28 day study by determination of oxygen demand in a closed respirometer.

Anox-Kaldnes AB, Lund, SE

2009

Test material:

Akoline PGPR, 100%

Results:

The test article is "readily biodegradable" according to the criteria specified in OECD guidelines for degradability testing.

Read across

Read across Similar chemical structures

Reference ID:

S168 - AnoxKaldnes 09-29-2

Test name:

Biodegradability

Method and laboratory:

OECD 301D

Test material:

Polyglycerol-3

Results:

Readily biodegradable

Read across

Read across Similar chemical structures

Reference ID:

S-352 ECHA, REACH registration 'Reaction mass of 3-[3-(2,3-dihydroxypropoxy)-2-hydroxypropoxylpropane-1,2-diol...',
<https://echa.europa.eu/registration-dossier/-/registered-dossier/13319>
Accessed 2020-08-18

Test name:

Biodegradability

Method and laboratory:

OECD 301F

Test material:

Polyglycerol-3 caprate/laurate
(read-across from polyglycerol-3 caprylate/caprylate)

Results:

Readily biodegradable

Read across

Read across Similar chemical structures

Reference ID:

S-353 ECHA, REACH registration 'Reaction products of decanoic acid and lauric acid with glycerol and polyglycerol',
<https://echa.europa.eu/registration-dossier/-/registered-dossier/25834>
Accessed 2020-08-18

Test name:

Biodegradability

Method and laboratory:

OECD 301F

Test material:

Polyglycerol-3 caprate

Results:

Readily biodegradable

Read across

Read across Similar chemical structures

Reference ID:

S-354 ECHA, REACH registration '1,2,3-propantriol, homopolymer, decanoate', CAS 74504-65-7

<https://echa.europa.eu/registration-dossier/-/registered-dossier/23621>
Accessed 2020-08-18

Test name:

Biodegradability

Method and laboratory:

OECD 301B

Test material:

Polyglycerol-3

Results:

Readily biodegradable

Read across

Read across Similar chemical structures

Reference ID:

S-355 ECHA, REACH registration '1,2,3-propantriol, homopolymer', CAS 25618-55-7
<https://echa.europa.eu/registration-dossier/-/registered-dossier/14285>
Accessed 2020-08-18

8.02 Accumulation

Polyglycerol and polyglycerol esters of fatty acids as well as relevant read-across substances are readily biodegradable. They are generally easily hydrolyzed to free fatty acids, polyglycerols and glycerol by aquatic and soil microorganisms. The fatty acids, polyglycerols and glycerol are easily metabolized by aquatic and soil microorganisms. Therefore the risk of environmental accumulation is regarded as minimal.

8.03 Aquatic toxicity

Test name:

Aquatic toxicity

Method and laboratory:

OECD 203 (Fish)

OECD 202 (Daphnia)

OECD 201 (Algae, *Pseudokirchneriella subcapitata*)

Test material:

Polyglycerol-3

Results:

LC50(96h, Fish): >1000 mg/l

EC50(48h, Daphnia): >1000 mg/l

NOEC(96h, Algae): 1000 mg/l

Read across

Read across Similar chemical structures

Reference ID:

S-352 ECHA, REACH registration 'Reaction mass of 3-[3-(2,3-dihydroxypropoxy)-2-hydroxypropoxy]propane-1,2-diol...',
<https://echa.europa.eu/registration-dossier/-/registered-dossier/13319>
Accessed 2020-08-18

Test name:

Aquatic toxicity

Method and laboratory:

OECD 203 (Fish)

OECD 202 (Daphnia)

OECD 201 (Algae, *Pseudokirchneriella subcapitata*)

Test material:

Polyglycerol-3 caprate/laurate

(read-across from polyglycerol-3 caprylate/caprylate)

Results:

LC50(96h, Fish): 32mg/l

EC50(48h, Daphnia): 32.1 mg/l

NOEC(72h, Algae): 50 mg/l

EC50(72h, Algae): 100 mg/l

Read across

Read across

Similar chemical structures

Reference ID:

S-353 ECHA, REACH registration 'Reaction products of decanoic acid and lauric acid with glycerol and polyglycerol',

<https://echa.europa.eu/registration-dossier/-/registered-dossier/25834>

Accessed 2020-08-18

Test name:

Aquatic toxicity

Method and laboratory:

QSAR VEGA SarPy/IFRMN 1.02 (Fish)

OECD 202 (Daphnia)

OECD 201 (Algae, *Pseudokirchneriella subcapitata*)

Test material:

Polyglycerol-3 caprate

Results:

LC50(QSAR Fish): 1-10 mg/l

EC50(48h, Daphnia): 15.1 mg/l

NOEC(72h, Algae): 10 mg/l

EC50(72h, Algae): 56.7 mg/l

Read across

Read across

Similar chemical structures

Reference ID:

S-354 ECHA, REACH registration '1,2,3-propantriol, homopolymer, decanoate', CAS 74504-65-7

<https://echa.europa.eu/registration-dossier/-/registered-dossier/23621>

Accessed 2020-08-18

Test name:

Aquatic toxicity

Method and laboratory:

OECD 203 (Fish)

OECD 202 (Daphnia)

OECD 201 (Algae, *Pseudokirchneriella subcapitata*)

Test material:

Polyglycerol-3

Results:

LC50(96h, Fish): >500 mg/l

EC50(48h, Daphnia): >1000 mg/l

EC50(72h, Algae): >1000 mg/l

Read across

Read across Similar chemical structures

Reference ID:

S-355 ECHA, REACH registration '1,2,3-propantriol, homopolymer', CAS 25618-55-7

<https://echa.europa.eu/registration-dossier/-/registered-dossier/14285>

Accessed 2020-08-18

8.04 Summary of ecotoxicity

The polyglycerol esters of fatty acids are readily biodegraded, do not accumulate in the environment and show low toxicity to aquatic organisms (fish, invertebrates and algae). It is therefore concluded that they pose minimal risk to the environment.

9.1 EU

9.1.1 Statement on EU Cosmetic Regulation EC 1223/2009

Latest statement, download "Statement on EU Cosmetic Regulation" at aakpersonalcare.com

9.1.2 EU Cosmetic Regulation EC 1223/2009, Annex II and III

Latest statement, download "Statement on EU Cosmetic Regulation" at aakpersonalcare.com

9.1.3 EU REACH 1907/2006

Latest statement, download "REACH Statement" at aakpersonalcare.com

9.1.4 EU SVHC (Substance of Very High Concern)

Latest statement, download "General Statement AAK Ingredients" at aakpersonalcare.com

9.2 Other country specific regulations:

9.2.1 US (California) Proposition 65

Latest statement, download "General Statement AAK Ingredients" at aakpersonalcare.com

9.2.2 China – NMPA

Latest statement, download "NMPA Statement" at aakpersonalcare.com

9.2.3 UK REACH

Latest statement, download "UK REACH Statements" at aakpersonalcare.com

9.2.4 Turkey - KKDIK

Latest statement, download "Turkey-KKDIK and SEA Statement" at aakpersonalcare.com

9.2.5 Australia - TGA

Latest statement, download "AAK PC Products and TGA status" at aakpersonalcare.com

9.3 Other non-Country specific regulatory issues

9.3.1 Animal testing

Latest statement, download "General Statement AAK Ingredients" at aakpersonalcare.com

9.3.2 Nano particles

Latest statement, download "General Statement AAK Ingredients" at aakpersonalcare.com

9.3.3 Nagoya Protocol / Biodiversity and Access Benefit Sharing regulation

Latest statement, download "General Statement AAK Ingredients" at aakpersonalcare.com

9.3.4 CITES

Latest statement, download "General Statement AAK Ingredients" at aakpersonalcare.com

9.3.5 CMR

Latest statement, download "General Statement AAK Ingredients" at aakpersonalcare.com

9.4 Inventory lists

Inventory lists relates to substances already existing in a specific market. The inventory list to the chemical legislation of the country or region. INCI labeling is not related to the chemical legislation. The nomenclature may differ between these two types of regulations hence the wording may change.

In the Table below, column 3:

- 1) Listed means:
 - a. The substance name and CAS number described as “AAK first choice name”, in section “1.1 Identification” is listed and not prohibited in the inventory list of the country.
- 2) Not listed, however CAS. No XXXXX-XX-X is listed and valid to be used.
 - a. The substance name and CAS number described as “AAK first choice name”, section “1.1 identification” is not found but instead the Cas XXXXX-XX-X mentions is listed as well as fits with the chemical description of the product, hence can be used instead.
- 3) No data:
 - a. AAK has not been able to find the substance in the inventory list.

EC (EU)	EC-inventory	Listed
TSCA (U.S.)	Toxic Substances Control Act	No
DSL (Canada)	Domestic Substances List	No but on the ICL list CAS 27321-72-8 is found.
AICS (Australia)	The Australian Inventory of Chemical Substances	No data
IECSC (China)	Inventory of Existing Chemical Substances Produced or Imported in China	No data
IECIC (China)	Inventory of Existing Cosmetic Ingredients in China	No data
ENCS (Japan)	Combined list of existing and notified chemical substances as the Japanese Existing and New Chemical Substances Inventory.	CAS 27321-72-8 is listed and valid to be used.
Japan	Japan Pharmacopoeia	Not found
KECI (South Korea)	Korea Existing Chemicals Inventory	No data
PICCS (Philippines)	Philippine Inventory of Chemicals and Chemical Substances	No data
NZIoC (New Zealand)	New Zealand Inventory of Chemicals	No data
NECI (Taiwan)	National Existing Chemical Inventory	Yes, stearic acid, monoester with triglycerol / Octadecanoic acid, monoester with triglycerol
Saudia Arabia	The Saudi Arabian Standards Organisation	No data
Malaysia	Chemicals Information Management System	No data
Mexico	Inventario Nacional de Sustancias Químicas	No data
Turkey		Yes. Local name:Stearik asit, trigliserollü monoester

10.1 Official standards

Standard	Conform	Monograph
EUR/Ph	n.a	
USP/NF	n.a	
JP	See inventory list 9.4	

10.2 Private standards

10.2.1 Ecocert, Cosmos or Natrue

Latest statement, download "General Statement AAK Ingredients" at aakpersonalcare.com

10.2.2 Vegan and Vegetarian claim

Latest statement, download "General Statement AAK Ingredients" at aakpersonalcare.com

10.2.3 Other

10.3 Other Statements

10.3.1 BSE/TSE statements:

Not available, please contact AAK for more information.

10.3.2 GMO statement

Not available, please contact AAK for more information.

10.3.3 Other:

No data

11. CERTIFICATES

11.1 Halal

Contact AAK for more information

.

11.2 Kosher

Contact AAK for more information

11.3 ISO 9001

Contact AAK for more information

11.4 EFFCI GMP

No data

11.5 Food Safety/ FSSC 22000

Contact AAK for more information

11.6 Other

No other available

12. PATENTS

12.1 Patents

No data.

TRANSPORTS AND HANDLING – Akoline PG7™

13.1 Transports

No data available

13.2 storage unopen package

Storage to fulfill shelf life:

Store in temperature below 20C or lower. Dark, dry and odour free condition in unopen packaging's.
See Product data sheet for more information.

Retest of batch:

Retest for prolonged shelf life is only possible after agreement with sales responsible.

13.3 Handling of product for use

13.3.1 Use of full package

Recommended melting temperature.

Bag is possible to use amount needed directly from the bag. Do not melt material directly in the bag remove and melt in a vessel.

During processing need to be heated to 65C to remove crystal memory.

13.3.2 Use of full package for partly use

Bag is possible to use partly directly from the bag. Do not melt material directly in the bag remove and melt in a vessel.

Reseal packaging and store in 20C or below

Note:

AAK's shelf life for ingredients that are unopened and stored according to the instructions given in the product information sheet. This guarantee is invalidated once the packaging is opened and the ingredients reheated. It is the user's responsibility to validate that a reheated material fulfills shelf life requirements in a formulation. See Product Data Sheet.

14. REFERENCES

14.1 References

No data

15. DISCLAIMER

15.1 Disclaimer

This document, or any answers or information provided herein by AAK, does not constitute a legally binding document of AAK. While the description designs, data and information contained herein are presented in good faith and believe to be accurate, it is provided for your guidance only. Because many factors may affect processing or application/use, we recommend that you make tests to determine the suitability of a product for your particular prior to use. It does not relieve our customers from obligation to perform a full inspection of the product upon delivery or any other obligation. No warranties of any kind either express or implied, including warranties of merchantability or fitness for a particular purpose are made regarding products described or designs, data or information set forth, or that the products design, data or information may be used without infringing the intellectual property right of others. In no case shall the descriptions, information, data or designs provided be considered a part of our terms and condition of sale.



Ship-to -

Analytical Certificate

Delivery	81303783 - 10
Print date	2023-03-21
Your reference	
Our reference	Angelique Mazur
Material	8691-805 Akoline PG7™
Your material no.	
Date of shipment	2023-03-21

Batch 4014310414 / **Quantity** 20 KG / **Prod. date** 2022-12-20
Inspection lot 3110574 / **Best before** 2024-12-19

Characteristic	Result	Lower Limit	Target	Upper Limit
Acid value(Syratal LB)				
Acid value	0,40 mg KOH/g			4,00
Saponification value(Försåpn Tal LB)				
Saponification value	137 mg KOH/g	130		145

Shelf life: 24 months from production date.

Quality Control Manager
AAK Sweden AB


This document is electronically produced, and valid without an AAK signature.

ZAO S11855 1

A Company in the AAK Group

AAK Sweden AB
SE-374 82 Karlshamn
SwedenPhone : +46(0)454 820 00
Website : www.aak.comBank : Skandinaviska Enskilda Banken
Bic/Swift : ESSESESS
Giro : 5430-5438
Acc. no. : 51181061768
IBAN : SE20 5000 0000 0511 8106 1768Org. no. : 556478-1796
VAT no. : SE556478179601
Approved for Swedish F-tax
Registered Office: Karlshamn
See Disclaimer

Flowchart Akoline Products



Flowchart
Akoline PG7